

GANIRELIX ACETATE 250 mcg



Description: A GnRH antagonist, which blocks GnRH receptors on the pituitary gonadotropin pathway. It induces a rapid, reversible suppression of gonadotropin secretion. Each package of Ganirelix Acetate contains a premixed, pre-filled syringe ready for subcutaneous injection.

How it works: Ganirelix Acetate is an injectable medication administered once daily subcutaneously. It is initiated a few days after FSH (follicle stimulating hormone) therapy and should be continued until the day of hCG administration. Ganirelix Acetate works immediately by suppressing LH and FSH secretion. This allows for eggs to reach the level of development needed for fertilization.

Potential side effects: Some women may experience headache, abdominal pain and nausea.

Storage: Store at room temperature (77°F); excursions permitted to (59-86°F). Protect from light.

Subcutaneous Administration of GANIRELIX ACETATE

- Cleanse your selected SUBCUTANEOUS injection site with an alcohol wipe and allow to dry.
- Remove cover from needle and invert the syringe so the needle is pointing upwards and tap the syringe to force any air bubbles to the top. Depress the plunger to expel the air.
- While holding the syringe with your dominant hand between your thumb and finger (as you would a pencil), pinch the skin (with your other hand) at the site you have selected.
- With a quick, dart-like motion, insert the entire needle (up to the hub of the syringe) at a 90° or a 45° angle and depress the plunger slowly and carefully.
- After the injection is complete, quickly pull out the needle. If bleeding, apply pressure to the injection site with a sterile gauze pad. If discomfort occurs after your injection, you may gently massage the injection site to alleviate your discomfort.
- Use Sharps waste container for needle and syringe.

