

CETROTIDE® 0.25 mg
CETROTIDE® 3.0 mg



Description: A GnRH antagonist, which blocks GnRH receptors on the pituitary gonadotropin pathway. It induces a rapid, reversible suppression of gonadotropin secretion. The vials of Cetrotide contain 0.25 mg and 3.0 mg of powdered medication packaged with prefilled syringes of diluent.


How it works: Cetrotide is an injectable medication administered subcutaneously. Cetrotide blocks the effect of the naturally occurring gonadotropin releasing hormone (GnRH). GnRH controls the release of LH that produces the release of an egg from the ovary. Cetrotide is given over a short period of time in the cycle when the LH surge can occur. It's blocking effect allows for eggs to reach the level of development needed for fertilization.

Potential side effects: Some women may experience headache, abdominal pain and nausea.

Storage: Cetrotide 0.25 mg - Store refrigerated (36-46°F). Store in original box. Use immediately after reconstitution/mixing.
Cetrotide 3.0 mg - Store at room temperature (68-77°F); excursions permitted to (59-86°F). Store in original box. Use immediately after reconstitution/mixing.

Subcutaneous Administration of CETROTIDE

- Pop off the plastic top of the vial of the Cetrotide.
- Swab the top of the vial with an alcohol pad to prevent contamination.
- Using the prefilled diluent syringe, slowly and cautiously deposit the diluent into the vial of powdered medication. Rotate the vial between your thumb and index finger until the solution is clear. Withdraw the newly mixed medication back into the syringe. Carefully recap and remove the mixing needle by twisting and replace it with a smaller subcutaneous injection needle provided.



- Invert the syringe so the needle is pointing upwards and tap the syringe to force any air bubbles to the top. Depress the plunger to expel the air.
- Cleanse your selected SUBCUTANEOUS injection site with an alcohol wipe and allow to dry.
- While holding the syringe with your dominant hand between your thumb and finger (as you would a pencil), pinch the skin (with your other hand) at the site you have selected.



- With a quick, dart-like motion, insert the entire needle (up to the hub of the syringe) at a 90° or a 45° angle and depress the plunger slowly and carefully.
- After the injection is complete, quickly pull out the needle. If bleeding, apply pressure to the injection site with a sterile gauze pad. If discomfort occurs after your injection, you may gently massage the injection site to alleviate your discomfort.
- Use Sharps waste container for needle and syringe.